

Variations in patterns of care for retroperitoneal sarcoma: a study of the Trans-Atlantic Australasian Retroperitoneal Sarcoma Working Group

- Brief overview:
 - Examines consistencies of use of various diagnostic, surgical, neo/adjuvant modalities in RPS
- Eligibility:
 - Enrolling patients into RESAR database between Feb 2017 – Jan 2020
- Data requirements outside of RESAR: None
- Contacts:
 - hayden.snow@petermac.org (study enquiries)
 - Marco.Fiore@istitutotumori.mi.it (RESAR DTA enquiries)
- Update:
 - Awaiting finalization of RESAR agreements – then can gain access to data and start analysis

Centers confirmed:

- Queen Elizabeth, Birmingham
- Peter Mac, Melbourne
- The Ottawa Hospital
- Ohio State University
- Leiden University Med Center
- European Institute of Oncology, Milan
- Dana-Farber/Brigham & Women's
- Mount Sinai/Princes Margaret, Toronto
- Humanitas University Hospital
- IOV, Padova
- Ludwig Maximilian University, Munich
- Sourasky Medical Centre, Tel-Aviv
- Hospital San Pau, Barcelona
- Peking University International Hospital, Beijing
- NCC, Tokyo
- Maria Sklodowska Curie, Warsaw
- Royal Marsden Hospital, London
- Institut Curie, Paris
- Winship Cancer Institute, Emory University
- Campus Biomedico, Roma
- Institute of Oncology, Ljubljana
- Samsung Medical Center, Seoul
- Candiolo Cancer Institute
- Netherlands Cancer Institute
- Mayo Clinic
- Istituto Nazionale dei Tumori
- University of Southern California, LA
- University of California, San Diego